

## HOMEWORK ASSIGNMENT 3, Math 253

1. Calculate the following limits, or discuss why they do not exist:

(a)  $\lim_{(x,y) \rightarrow 0} \frac{y}{x^2 + y^2}$

(b)  $\lim_{(x,y) \rightarrow 0} \frac{y^3}{x^2 + y^2}$  [hint:  $|y^2| \leq |x^2 + y^2|$ ]

2. For each of the following functions, give its domain and calculate  $\frac{\partial f}{\partial x}$  and  $\frac{\partial f}{\partial y}$ :

(a)  $f(x, y) = e^{3x} \cos(3y)$

(b)  $f(x, y) = \ln(1 + xy^2)$

(c)  $f(x, y) = \frac{y}{x^2 + y^2}$

(d)  $f(x, y) = x^y$

(e)  $f(x, y) = \cosh(x) \cos(y)$

(f)  $f(x, y) = x^3 \arcsin y^2$

3. Which of the above functions satisfies the Laplace equation:  $f_{xx} + f_{yy} = 0$ ?

4. Give an equation for the tangent plane to the graph of  $f(x, y) = \frac{y}{x^2 + y^3}$  at the point  $(0, 1, 1)$ . What is the normal vector at that point?

5. Find the coordinates of all points at which the surface with the following equation has a horizontal tangent plane:  $z = x^4 - 4xy^3 + 6y^2 - 2$ .

6. The equation  $x^3y^4 + xz^2 - yz^3 = 1$  defines a surface which passes through the point  $(1, 1, 1)$ . Find  $\frac{\partial z}{\partial x}$  and  $\frac{\partial z}{\partial y}$  at that point.

7. The radius and height of a right circular-conical tank are measured and found to be 25 *ft* and 21 *ft* respectively. Each measurement is accurate to within 0.5 *in*. By about how much can the calculated volume of the tank be in error?

8. Write an appropriate version of the chain rule for  $\frac{\partial z}{\partial u}$ , if  $z = g(x, y)$ ,  $y = f(x)$  and  $x = h(u, v)$ .

9. Use two different methods to calculate  $\frac{\partial z}{\partial x}$  given that

$$z = \arctan(u/v), \quad u = 2x + y, \quad v = 3x - y.$$

10. Calculate  $\frac{\partial}{\partial x} f(y^2, x^2)$ , assuming  $f$  has continuous partial derivatives.

## SOLUTIONS TO HOMEWORK ASSIGNMENT 3, Math 253

1. Calculate the following limits, or discuss why they do not exist:

(a)  $\lim_{(x,y) \rightarrow 0} \frac{y}{x^2 + y^2}$

**Solution:** The limit does not exist. If  $(x, y)$  approaches  $(0, 0)$  along the  $x$ -axis, we have  $y = 0, x \neq 0$  and the expression is identically zero, and so the limit in that direction is zero. On the other hand, approaching  $(0, 0)$  along the  $y$ -axis, the expression becomes  $1/y$ , which increases without bound as  $y \rightarrow 0$ , so the limit cannot exist (even as  $\infty$ ).

(b)  $\lim_{(x,y) \rightarrow 0} \frac{y^3}{x^2 + y^2}$  [hint:  $|y^2| \leq |x^2 + y^2|$ ]

**Solution:** Note that  $0 \leq \left| \frac{y^3}{x^2 + y^2} \right| = \frac{|y^2|}{|x^2 + y^2|} |y| \leq |y|$ , the latter inequality following from the hint. Since  $|y| \rightarrow 0$ , the same must be true of  $\left| \frac{y^3}{x^2 + y^2} \right|$ , so  $\lim_{(x,y) \rightarrow 0} \frac{y^3}{x^2 + y^2} = 0$ .

2. For each of the following functions, give its domain and calculate  $\frac{\partial f}{\partial x}$  and  $\frac{\partial f}{\partial y}$ :

(a)  $f(x, y) = e^{3x} \cos(3y)$

**Solution:** The domain is all of  $\mathbb{R}^2$ .

$$\frac{\partial f}{\partial x} = 3e^{3x} \cos(3y) \text{ and } \frac{\partial f}{\partial y} = -3e^{3x} \sin(3y).$$

(b)  $f(x, y) = \ln(1 + xy^2)$

**Solution:** Since the logarithm requires a positive input, we must have  $xy^2 > -1$ , or equivalently  $x > -\frac{1}{y^2}$ . The graph of  $x > -\frac{1}{y^2}$  has 2 branches, one in the third quadrant and one in the fourth. The domain of definition is everything between these graphs together the set of all points  $(x, y)$  such that  $x \geq 0$ , i.e.

$$\left\{ (x, y) \mid x < 0, -\sqrt{-1/x} < y < \sqrt{-1/x} \right\} \cup \left\{ (x, y) \mid x \geq 0 \right\}$$

$$\frac{\partial f}{\partial x} = y^2/(1 + xy^2) \text{ and } \frac{\partial f}{\partial y} = 2xy/(1 + xy^2)$$

(c)  $f(x, y) = \frac{y}{x^2 + y^2}$

**Solution:** The domain is  $\mathbb{R} \setminus (0, 0)$ .  $\frac{\partial f}{\partial x} = -2xy/(x^2 + y^2)^2$  and  $\frac{\partial f}{\partial y} = (x^2 - y^2)/(x^2 + y^2)^2$

(d)  $f(x, y) = x^y$

**Solution:** The standard definition is  $x^y = e^{(y \ln x)}$ , which requires  $x > 0$ , so the domain is the open right half-plane.  $\frac{\partial f}{\partial x} = yx^{y-1}$  and  $\frac{\partial f}{\partial y} = x^y \ln x$ .

(e)  $f(x, y) = \cosh(x) \cos(y)$

**Solution:** The domain is all of  $\mathbb{R}^2$ .  $\frac{\partial f}{\partial x} = \sinh(x) \cos(y)$  and  $\frac{\partial f}{\partial y} = -\cosh(x) \sin(y)$

(f)  $f(x, y) = x^3 \arcsin y^2$

**Solution:** The arcsin requires the input  $y^2$  between  $-1$  and  $+1$ , which is equivalent to  $-1 \leq y \leq 1$ . So the range is the infinite horizontal strip between the lines  $y = -1$  and  $y = 1$ .  $\frac{\partial f}{\partial x} = 3x^2 \arcsin y^2$  and  $\frac{\partial f}{\partial y} = 2x^3 y / \sqrt{1 - y^4}$ .

3. Which of the above functions satisfies the Laplace equation:  $f_{xx} + f_{yy} = 0$ ?

**Solution:** (a), (c) and (e).

For (a):  $f_{xx} = 9e^x \cos 3y = -f_{yy}$ .

For (c):  $f_{xx} = (6x^2y - 2y^3)/(x^2 + y^2)^3 = -f_{yy}$ .

For (e):  $f_{xx} = \cosh(x) \cos(y) = -f_{yy}$ .

4. Give an equation for the tangent plane to the graph of  $f(x, y) = \frac{y}{x^2 + y^3}$  at the point  $(0, 1, 1)$ . What is the normal vector at that point?

**Solution:** Since  $f_x(0, 1) = 0$  and  $f_y(0, 1) = -2$ , the tangent plane has equation  $z = 1 + (-2)(y - 1)$  or  $2y + z = 3$ . A normal vector at  $(0, 1, 1)$  is  $\langle 0, 2, 1 \rangle$ .

5. Find the coordinates of all points at which the surface with the following equation has a horizontal tangent plane:  $z = x^4 - 4xy^3 + 6y^2 - 2$ .

**Solution:** There is a horizontal tangent plane when  $\frac{\partial z}{\partial x} = \frac{\partial z}{\partial y} = 0$ . This gives  $4x^3 - 4y^3 = 0$  and  $-12xy^2 + 12y = 0$ . The first equation implies  $x = y$ . The second can be rewritten  $y(1 - y^2) = 0$ , and so  $y = 0$  or  $y = \pm 1$ . Thus there are three points at which the surface has a horizontal tangent plane:  $(0, 0, -2)$ ;  $(1, 1, 1)$  and  $(-1, -1, 1)$ .

6. The equation  $x^3y^4 + xz^2 - yz^3 = 1$  defines a surface which passes through the point  $(1, 1, 1)$ . Find  $\frac{\partial z}{\partial x}$  and  $\frac{\partial z}{\partial y}$  at that point.

**Solution:** We can find the partials by implicit differentiation. Holding  $y$  fixed and taking the derivative with respect to  $x$  (regarding  $z$  as function of  $x$ ) gives

$$3x^2y^4 + z^2 + 2xz \frac{\partial z}{\partial x} - 3yz^2 \frac{\partial z}{\partial x} = 0.$$

At the point  $(1, 1, 1)$  this becomes  $3 + 1 + 2\frac{\partial z}{\partial x} - 3\frac{\partial z}{\partial x} = 0$ , so

$\frac{\partial z}{\partial x}(1, 1) = 4$ . Similarly, holding  $x$  fixed and taking the  $y$ -derivative gives

$$4x^3y^3 + 2xz\frac{\partial z}{\partial y} - z^3 - 3yz^2\frac{\partial z}{\partial y} = 0,$$

which becomes  $4 + 2\frac{\partial z}{\partial y} - 1 - 3\frac{\partial z}{\partial y} = 0$ , and we find  $\frac{\partial z}{\partial y}(1, 1) = 3$ .

7. The radius and height of a right circular-conical tank are measured and found to be  $25 \text{ ft}$  and  $21 \text{ ft}$  respectively. Each measurement is accurate to within  $0.5 \text{ in.}$  By about how much can the calculated volume of the tank be in error?

**Solution:** The volume of a right circular cone of radius  $r$  and height  $h$  is  $V = \pi r^2 h/3$ . Then

$$dV = \frac{2\pi}{3} r h dr + \frac{\pi}{3} r^2 dh = \frac{2\pi}{3} \times 25 \times 21 \times \frac{1}{24} + \frac{\pi}{3} (25)^2 \frac{1}{24} = 1675\pi/24 \approx 73(\text{ft}^3).$$

8. Write an appropriate version of the chain rule for  $\frac{\partial z}{\partial u}$ , if  $z = g(x, y)$ ,  $y = f(x)$  and  $x = h(u, v)$ .

**Solution:**  $\frac{\partial z}{\partial u} = g_1 h_1 + g_2 f' h_2$ , or in Leibniz notation  $\frac{\partial z}{\partial u} = \frac{\partial g}{\partial x} \frac{\partial h}{\partial u} + \frac{\partial g}{\partial y} \frac{df}{dx} \frac{\partial h}{\partial u}$ .

9. Use two different methods to calculate  $\frac{\partial z}{\partial x}$  given that

$$z = \arctan(u/v), \quad u = 2x + y, \quad v = 3x - y.$$

**Solution:** Using either direct substitution or the chain rule results in the answer

$$\frac{\partial z}{\partial x} = \frac{-5y}{13x^2 - 2xy + 2y^2}.$$

10. Calculate  $\frac{\partial}{\partial x} f(y^2, x^2)$ , assuming  $f$  has continuous partial derivatives.

**Solution:**  $2x f_2(y^2, x^2)$ .