

# PLANE-FILLING CURVES OF SMALL DEGREE OVER FINITE FIELDS

SHAMIL ASGARLI AND DRAGOS GHIOCA

ABSTRACT. A plane curve  $C$  in  $\mathbb{P}^2$  defined over  $\mathbb{F}_q$  is called plane-filling if  $C$  contains every  $\mathbb{F}_q$ -point of  $\mathbb{P}^2$ . Homma and Kim, building on the work of Tallini, proved that the minimum degree of a smooth plane-filling curve is  $q + 2$ . We study smooth plane-filling curves of degree  $q + 3$  and higher.

## 1. INTRODUCTION

The study of space-filling curves in  $\mathbb{R}^2$  starts with the work of Peano [Pea90] in the 19th century. About 100 years later, Nick Katz [Kat99] studied space-filling curves over finite fields and raised open questions about their existence. One version of Katz's question was the following. Given a smooth algebraic variety  $X$  over a finite field  $\mathbb{F}_q$ , does there always exist a *smooth* curve  $C \subset X$  such that  $C(\mathbb{F}_q) = X(\mathbb{F}_q)$ ? In other words, is it possible to pass through all of the (finitely many)  $\mathbb{F}_q$ -points of  $X$  using a smooth curve? Gabber [Gab01] and Poonen [Poo04] independently answered this question in the affirmative.

We will consider the special case when  $X = \mathbb{P}^2$ . We say that a curve  $C \subset \mathbb{P}^2$  is *plane-filling* if  $C(\mathbb{F}_q) = \mathbb{P}^2(\mathbb{F}_q)$ . Equivalently,  $C$  is a plane-filling curve if  $\#C(\mathbb{F}_q) = q^2 + q + 1$ . In a natural sense, plane-filling curves are extremal. There are other classes of extremal curves with respect to the set of  $\mathbb{F}_q$ -points, including blocking curves [AGY23] and tangent-filling curves [AG23].

From Poonen's work [Poo04], we know that there exist smooth plane-filling curves of degree  $d$  over  $\mathbb{F}_q$  whenever  $d$  is sufficiently large with respect to  $q$ . It is natural to ask for the minimum degree of a smooth plane-filling curve over  $\mathbb{F}_q$ . Homma and Kim [HK13] proved that the minimum degree is  $q + 2$ . More precisely, by building on the work of Tallini [Tal61a, Tal61b], they showed that a plane-filling curve of the form

$$(ax + by + cz)(x^q y - xy^q) + y(y^q z - yz^q) + z(z^q x - zx^q) = 0$$

is smooth if and only if the polynomial  $t^3 - (ct^2 + bt + a) \in \mathbb{F}_q[t]$  has no  $\mathbb{F}_q$ -roots. In a sequel paper [Hom20], Homma investigated further properties of plane-filling curves of degree  $q + 2$ . The automorphism group of these special curves was studied by Duran Cunha [DC18]. As another direction, Homma and Kim [HK23] investigated space-filling curves in  $\mathbb{P}^1 \times \mathbb{P}^1$ .

In this paper, we investigate the existence of smooth plane-filling curves of degree  $q + 3$  and higher. The guiding question for our paper is the following.

**Question 1.1.** Let  $q$  be a prime power. Does there exist a smooth plane-filling curve of degree  $q + 3$  defined over  $\mathbb{F}_q$ ?

The three binomials  $x^q y - xy^q$ ,  $y^q z - yz^q$ , and  $z^q x - zx^q$  generate the ideal of polynomials defining plane-filling curves; see [HK13, Proposition 2.1] for proof of this assertion. Thus, any

---

2020 *Mathematics Subject Classification.* Primary: 14G15, 14H50; Secondary: 11G20, 14G05.

*Key words and phrases.* Plane curve, space-filling curve, smooth curve, finite field.

plane-filling curve of degree  $q + 3$  must necessarily be defined by

$$Q_1(x, y, z) \cdot (x^q y - x y^q) + Q_2(x, y, z) \cdot (y^q z - y z^q) + Q_3(x, y, z) \cdot (z^q x - z x^q) = 0$$

for some homogeneous quadratic polynomials  $Q_1, Q_2, Q_3 \in \mathbb{F}_q[x, y, z]$ . The difficulty is finding suitable  $Q_1, Q_2, Q_3$  for which the corresponding curve is smooth.

Our first result gives a necessary and sufficient condition for the plane-filling curve  $C_k$  to be smooth at all the  $\mathbb{F}_q$ -points.

**Theorem 1.2.** *For each  $k \in \mathbb{F}_q$ , consider the plane-filling curve  $C_k$  defined by*

$$x^2(x^q y - x y^q) + y^2(y^q z - y z^q) + (z^2 + kx^2)(z^q x - z x^q) = 0. \quad (1)$$

*Then  $C_k$  is smooth at every  $\mathbb{F}_q$ -point of  $\mathbb{P}^2$  if and only if the polynomial  $x^7 + kx^3 - 1$  has no zeros in  $\mathbb{F}_q$ .*

To ensure that the previous theorem is not vacuous, we need to show that there exists some  $k \in \mathbb{F}_q$  such that  $x^7 + kx^3 - 1$  has no zeros in  $\mathbb{F}_q$ .

**Proposition 1.3.** *There exists a value  $k \in \mathbb{F}_q$  such that  $x^7 + kx^3 - 1 \in \mathbb{F}_q[x]$  has no zeros in  $\mathbb{F}_q$ .*

*Proof.* When  $x = 0$ , there is no  $k \in \mathbb{F}_q$  such that  $x^7 + kx^3 - 1 = 0$ . For each  $x \in \mathbb{F}_q^*$ , there is a unique value of  $k \in \mathbb{F}_q$  such that  $x^7 + kx^3 - 1 = 0$ . Thus, there are at most  $q - 1$  values of  $k \in \mathbb{F}_q$  such that the polynomial  $x^7 + kx^3 - 1$  has a zero in  $\mathbb{F}_q$ .  $\square$

The next result improves Proposition 1.3.

**Theorem 1.4.** *There exist at least  $\frac{q}{6} - 1 - \frac{28}{3}\sqrt{q}$  many values of  $k \in \mathbb{F}_q$  such that  $x^7 + kx^3 - 1 \in \mathbb{F}_q[x]$  has no zeros in  $\mathbb{F}_q$ .*

Note that Theorem 1.2 and Proposition 1.3 together yields that for each odd  $q$ , there exists at least one value  $k \in \mathbb{F}_q$  for which the corresponding curve  $C_k$  has no singular  $\mathbb{F}_q$ -points. Furthermore, we expect that the curves in Theorem 1.2 are smooth if and only if they are smooth at all their  $\mathbb{F}_q$ -points. Our main conjecture below restates this prediction.

**Conjecture 1.5.** *Suppose  $q$  is odd. The plane-filling curve  $C_k$  defined by (1) is smooth if and only if the polynomial  $x^7 + kx^3 - 1$  has no zeros in  $\mathbb{F}_q$ .*

We have verified Conjecture 1.5 using Macaulay2 [GS] for all odd prime powers  $q < 200$ . When  $q = 2^m$  is even, the curve  $C_k$  defined by (1) turns out to be singular (for every  $k \in \mathbb{F}_q$ ). As a replacement, we consider another curve  $D_k$  in this case:

$$x^2(x^q y - x y^q) + y^2(y^q z - y z^q) + (z^2 + kxy)(z^q x - z x^q) = 0. \quad (2)$$

We make a similar conjecture regarding the smoothness of the curves  $D_k$ .

**Conjecture 1.6.** *Suppose  $q$  is even. The plane-filling curve  $D_k$  defined by (2) is smooth if and only if the polynomial  $x^7 + kx^5 + 1$  has no zeros in  $\mathbb{F}_q$ .*

The polynomial  $x^7 + kx^5 + 1$  featured above is prominent because one can show, similar to Theorem 1.2, that a plane-filling curve  $D_k$  is smooth at all of its  $\mathbb{F}_q$ -points (when  $q$  is even) if and only if  $x^7 + kx^5 + 1$  has no  $\mathbb{F}_q$ -roots. We have verified Conjecture 1.6 using Macaulay2 [GS] for  $q = 2^m$  when  $1 \leq m \leq 9$ .

We prove the following as partial progress towards Conjecture 1.5.

**Theorem 1.7.** *Suppose  $q$  is odd. There exists a suitable choice of  $k \in \mathbb{F}_q$  such that the plane-filling curve  $C_k$  defined by (1) is smooth at all  $\mathbb{F}_{q^2}$ -points.*

A similar argument as the one employed in Theorem 1.7 yields an analogous result when  $q$  is even, and the curve  $C_k$  is replaced by  $D_k$ .

To prove Theorem 1.7, we will prove that any plane-filling curve of degree  $q+3$  which is smooth at  $\mathbb{F}_q$ -points and has no  $\mathbb{F}_q$ -linear component must be smooth at each of its  $\mathbb{F}_{q^2}$ -points.

We also investigate plane-filling curves of degree  $q+r+1$  where  $r \geq 2$  is arbitrary.

**Theorem 1.8.** *For each  $k \in \mathbb{F}_q$ , consider the plane-filling curve  $C_{k,r}$  defined by*

$$x^r(x^q y - xy^q) + y^r(y^q z - yz^q) + (z^r + kx^r)(z^q x - zx^q) = 0.$$

*Then  $C_{k,r}$  is smooth at every  $\mathbb{F}_q$ -point of  $\mathbb{P}^2$  if and only if the polynomial  $x^{r^2+r+1} + kx^{r+1} - 1 = 0$  has no zeros in  $\mathbb{F}_q$ .*

**Structure of the paper.** In Section 2, we prove Theorem 1.4. We devote Section 3 to Theorem 1.7, and Section 4 to Theorem 1.8.

## 2. PROOF OF THEOREM 1.4

We begin this section by noting that Theorem 1.2 is a special case of Theorem 1.8 which will be proven in Section 4. Our Theorem 1.2 provides a criterion that tests whether the plane-filling curve  $C_k$  defined by (1) is smooth at every  $\mathbb{F}_q$ -point.

The following technical result will be employed in our proof of Theorem 1.4.

**Lemma 2.1.** *The polynomial  $x^3 y^3(x+y)(x^2+y^2) + (x^2+xy+y^2)$  is irreducible in  $\overline{\mathbb{F}_q}[x, y]$ .*

*Proof.* The proof employs a technique seen in Eisenstein's criterion. First, suppose  $p = \text{char}(\mathbb{F}_q) \neq 3$ . Assume, to the contrary, that  $f(x, y) := x^3 y^3(x+y)(x^2+y^2) + (x^2+xy+y^2)$  is reducible over the algebraic closure  $\overline{\mathbb{F}_q}$ . Write  $f(x, y) = g(x, y) \cdot h(x, y)$ , and express

$$\begin{aligned} g(x, y) &= g_m(x, y) + g_{m+1}(x, y) + \cdots + g_s(x, y) \\ h(x, y) &= h_n(x, y) + h_{n+1}(x, y) + \cdots + h_t(x, y) \end{aligned}$$

where  $g_i(x, y)$  and  $h_j(x, y)$  are homogeneous of degree  $i$  and  $j$ , respectively, for  $m \leq i \leq s$  and  $n \leq j \leq t$ . From  $f(x, y) = g(x, y) \cdot h(x, y)$ , we see that

$$\begin{cases} g_m h_n = x^2 + xy + y^2 \\ g_s h_t = x^3 y^3(x+y)(x^2+y^2) \\ \sum_{i+j=k} h_i g_j = 0 \text{ for } 2 < k < 9 \end{cases}$$

Since the characteristic  $p \neq 3$ , the polynomial  $x^2 + xy + y^2$  factors into distinct linear factors in  $\overline{\mathbb{F}_q}[x, y]$ . Let  $x + \lambda y$  be one of those linear factors with  $\lambda \in \overline{\mathbb{F}_q}$ . Then  $x^2 + xy + y^2$  is divisible by  $x + \lambda y$  but not by  $(x + \lambda y)^2$ . Thus, exactly one of  $g_m$  or  $h_n$  is divisible by  $x + \lambda y$ . Without loss of generality, assume  $x + \lambda y$  divides  $g_m$ , and not  $h_n$ . Then using  $\sum_{i+j=k} h_i g_j = 0$  for  $2 < k < 9$ , we inductively see that  $x + \lambda y$  divides  $g_j$  for each  $m \leq j \leq s$ . In particular,  $x + \lambda y$  divides  $g_s h_t$ . This is a contradiction because  $x + \lambda y$  does not divide  $x^3 y^3(x+y)(x^2+y^2)$ . Indeed,  $x^2 + xy + y^2$  and  $x^3 y^3(x+y)(x^2+y^2)$  are relatively prime.

When  $p = 3$ , a similar argument works from the other end of the polynomial: the leading term  $x^3 y^3(x+y)(x^2+y^2)$  is divisible by  $x+y$  but not by  $(x+y)^2$ . We deduce that  $f(x, y)$  is irreducible over  $\overline{\mathbb{F}_q}$  for every prime power  $q$ .  $\square$

*Proof of Theorem 1.4.* Our goal is to give a lower bound on the number of  $k \in \mathbb{F}_q$  such that the polynomial  $x^7 + kx^3 - 1$  has no roots in  $\mathbb{F}_q$ . As  $x$  ranges in  $\mathbb{F}_q^*$  (note that there is no  $k \in \mathbb{F}_q$  for which  $x = 0$  would be a root of  $x^7 + kx^3 - 1$ ), the number of “bad” choices of  $k$  are parametrized by  $\frac{1-x^7}{x^3}$ . We will show that there are many choices of  $x$  and  $y$  such that  $\frac{1-x^7}{x^3}$  and  $\frac{1-y^7}{y^3}$  give rise to the same value of  $k$ . Setting these expressions equal to each other, we obtain the following.

$$\frac{1-x^7}{x^3} = \frac{1-y^7}{y^3} \Rightarrow x^7y^3 - y^3 = y^7x^3 - x^3$$

After rearranging and dividing both sides by  $x - y$ , we obtain an affine curve  $\mathcal{C} \subset \mathbb{A}^2$  defined by

$$x^3y^3(x+y)(x^2+y^2) + x^2 + xy + y^2 = 0,$$

for  $x, y \in \mathbb{F}_q^*$  and  $x \neq y$ . Let  $G$  be a graph whose vertex set is  $\mathbb{F}_q^*$ , and there is an edge between  $x$  and  $y$  if  $(x, y)$  lies on the affine curve  $\mathcal{C}$ . We consider undirected edges, so the pairs  $(x, y)$  and  $(y, x)$  correspond to the same edge.

**Claim 1.** The number of edges of  $G$  is at least  $\frac{q}{2} - 6 - 28\sqrt{q}$ .

Let  $\tilde{\mathcal{C}} \subset \mathbb{P}^2$  be the projectivization of  $\mathcal{C}$ . By Lemma 2.1, the curve  $\tilde{\mathcal{C}}$  is geometrically irreducible. By Hasse-Weil inequality for geometrically irreducible curves [AP96, Corollary 2.5],  $\#\tilde{\mathcal{C}}(\mathbb{F}_q) \geq q + 1 - 56\sqrt{q}$ . Since the line at infinity  $z = 0$  can contain at most 5 distinct  $\mathbb{F}_q$ -points, we have  $\#C(\mathbb{F}_q) \geq q - 4 - 56\sqrt{q}$ ; furthermore, we exclude the points for which  $xy = 0$  and there is only one such point  $[0 : 0 : 1] \in \tilde{\mathcal{C}}$ . We also need to rule out the points on the diagonal, namely  $x = y$ ; in this case,  $4x^9 + 3x^2 = 0$  which contributes at most 7 additional points with  $x \neq 0$ . Thus, the number of  $(x, y) \in C(\mathbb{F}_q)$  with  $x \neq y$  is at least  $q - 12 - 56\sqrt{q}$ . The claim follows since the edges are undirected.

**Claim 2.** Every connected component of  $G$  is a complete graph  $K_n$  where  $n \in \{1, 2, 3, 4, 5, 6\}$ .

If  $(x, y)$  and  $(x, z)$  are both edges of  $G$ , then  $\frac{1-x^7}{x^3} = \frac{1-y^7}{y^3}$  and  $\frac{1-x^7}{x^3} = \frac{1-z^7}{z^3}$ . Consequently,  $\frac{1-y^7}{y^3} = \frac{1-z^7}{z^3}$  and  $(y, z)$  lies on the curve  $\mathcal{C}$ , so  $(y, z)$  is an edge in  $G$  too. Thus, each connected component of  $G$  is a clique. In addition, from the equation of  $\mathcal{C}$ , the degree of each vertex  $x \in G$  is at most 6.

For each  $1 \leq i \leq 6$ , let  $m_i$  denote the number of cliques of size  $i$  in  $G$ . Counting the number of edges in  $G$  leads to the following equality.

$$\#E(G) = \sum_{i=1}^6 \frac{i(i-1)}{2} \cdot m_i.$$

Each clique of size  $i$  in  $G$  increases the number of “good” values of  $k$  by an additive factor of  $i - 1$  because each clique corresponds to one “bad” value of  $k$ , i.e., a value  $k \in \mathbb{F}_q$  for which the equation  $x^7 + kx^3 - 1 = 0$  is solvable for some  $x \in \mathbb{F}_q$ . More precisely,

$$\begin{aligned} & \#\{k \in \mathbb{F}_q \mid x^7 + kx^3 - 1 \text{ has no zeros in } \mathbb{F}_q\} \\ &= q - \sum_{i=1}^6 m_i \\ &= 1 + (q - 1) - \sum_{i=1}^6 m_i \end{aligned}$$

$$\begin{aligned}
&= 1 + \sum_{i=1}^6 i \cdot m_i - \sum_{i=1}^6 m_i \\
&= 1 + \sum_{i=1}^6 (i-1) \cdot m_i \\
&\geq 1 + \frac{1}{3} \sum_{i=1}^6 \frac{(i-1)i}{2} \cdot m_i \geq 1 + \frac{1}{3} \#E(G) \geq 1 + \frac{1}{3} \left( \frac{q}{2} - 6 - 28\sqrt{q} \right)
\end{aligned}$$

as desired.  $\square$

### 3. SMOOTHNESS AT $\mathbb{F}_{q^2}$ -POINTS

In this section, we show that a plane-filling curve  $C$  of degree  $q + 3$  has the following special property: being smooth at  $\mathbb{F}_q$ -points implies being smooth at  $\mathbb{F}_{q^2}$ -points under a mild condition.

**Proposition 3.1.** *Suppose  $C$  is a plane-filling curve of degree  $q + 3$  such that*

- (i) *The curve  $C$  is smooth at all the  $\mathbb{F}_q$ -points.*
- (ii) *The curve  $C$  has no  $\mathbb{F}_q$ -linear component.*

*Then  $C$  is smooth at each  $\mathbb{F}_{q^2}$ -point.*

*Proof.* Assume, to the contrary, that  $C$  is singular at some  $\mathbb{F}_{q^2}$ -point  $Q$ . Then  $Q$  is not an  $\mathbb{F}_q$ -point due to the hypothesis (i). Let  $Q^\sigma$  denote the Galois conjugate of  $Q$  under the Frobenius automorphism. More explicitly, if  $Q = [x : y : z] \in \mathbb{P}^2$ , then  $Q^\sigma = [x^q : y^q : z^q]$ . Note that  $Q^\sigma$  is also contained in  $C$  (since  $C$  is defined over  $\mathbb{F}_q$ ). Moreover,  $Q^\sigma$  is also a singular point of  $C$ .

Consider the line  $L$  joining  $Q$  and  $Q^\sigma$ , which is an  $\mathbb{F}_q$ -line by Galois theory. By hypothesis (ii), the line  $L$  must intersect  $C$  in exactly  $q + 3$  points (counted with multiplicity). However,  $L$  already contains  $q + 1$  distinct  $\mathbb{F}_q$ -points of  $C$  (because  $C$  is plane-filling), and passes through the two singular points  $Q$  and  $Q^\sigma$ , each contributing intersection multiplicity at least 2. Thus, the total intersection multiplicity between  $L$  and  $C$  is at least  $(q + 1) + 2 + 2 = q + 5$ , a contradiction.  $\square$

**Remark 3.2.** We can weaken the hypothesis of Proposition 3.1 by replacing the condition  $\deg(C) = q + 3$  with  $\deg(C) \leq q + 4$ . Indeed, the same proof works verbatim.

Next, we show that the plane-filling curves  $C_k$  of degree  $q + 3$  considered in equation (1) indeed satisfy condition (ii) when  $q$  is odd.

**Proposition 3.3.** *The curve  $C_k$  defined by (1) has no  $\mathbb{F}_q$ -linear components when  $q$  is odd.*

*Proof.* There are three types of  $\mathbb{F}_q$ -lines in  $\mathbb{P}^2$ .

**Type I.** The line  $L$  is given by  $z = 0$ .

The curve  $C_k$  meets the line  $\{z = 0\}$  at finitely many points determined by  $x^2(x^q y - xy^q) = 0$ . In particular,  $\{z = 0\}$  is not a component of  $C$ .

**Type II.** The line  $L$  is given by  $x = az$  for some  $a \in \mathbb{F}_q$ .

The curve  $C_k$  meets the line  $\{x = az\}$  at finitely many points determined by

$$(az)^2((az)^q y - (az)y^q) + y^2(y^q z - yz^q) + (z^2 + k(az)^2)(z^q(az) - z(az)^q) = 0.$$

After simplifying and using  $a^q = a$ , the last term cancels and we obtain:

$$a^3 z^{q+2} y - a^3 z^3 y^q + y^{q+2} z - y^3 z^q = 0$$

In particular,  $\{x = az\}$  is not a component of  $C$ .

**Type III.** The line  $L$  is given by  $y = ax + bz$  for some  $a, b \in \mathbb{F}_q$ .

If  $a = 0$  or  $b = 0$ , then  $y = bz$  or  $y = ax$ , and the analysis is very similar to the previous case. We will assume that  $a \neq 0$  and  $b \neq 0$ . We substitute  $y = ax + bz$  into the equation (1) and collect terms to obtain:

$$\begin{aligned} & (b + a^3 - k)x^{q+2}z + (2a^2b)x^{q+1}z^2 + (b^2a - 1)x^qz^3 + \\ & (-b - a^3 + k)x^3z^q + (-2ab)x^2z^{q+1} + (-ab^2 + 1)xz^{q+2} = 0 \end{aligned}$$

The coefficient of  $x^{q+1}z^2$  is  $2a^2b$ , which is nonzero since  $q$  is odd (so  $2 \neq 0$ ),  $a \neq 0$  and  $b \neq 0$ . Thus,  $L$  is not a component of  $C_k$ .  $\square$

We are now in a position to prove Theorem 1.7 on the existence of  $k \in \mathbb{F}_q$  such that the plane-filling curve  $C_k$  is smooth at all its  $\mathbb{F}_{q^2}$ -points.

*Proof of Theorem 1.7.* The result follows immediately from Proposition 1.3, Proposition 3.1, and Proposition 3.3.  $\square$

#### 4. HIGHER DEGREE PLANE-FILLING CURVES

We begin by establishing Theorem 1.8, which provides a necessary and sufficient condition for the plane-filling curve  $C_{k,r}$  to be smooth at all the  $\mathbb{F}_q$ -points.

*Proof of Theorem 1.8.* We consider the curve  $C_{k,r}$  given by the equation:

$$x^r \cdot (x^q y - xy^q) + y^r \cdot (y^q z - yz^q) + (z^r + kx^r) \cdot (z^q x - zx^q) = 0. \quad (3)$$

We analyze the singular locus of  $C_{k,r}$  and get the equations:

$$rx^{r-1} \cdot (x^q y - xy^q) + x^r \cdot (-y^q) + kr^{r-1} \cdot (z^q x - zx^q) + (z^r + kx^r) \cdot z^q = 0 \quad (4)$$

$$x^r \cdot x^q + ry^{r-1} \cdot (y^q z - yz^q) + y^r \cdot (-z^q) = 0 \quad (5)$$

$$y^r \cdot y^q + rz^{r-1} \cdot (z^q x - zx^q) + (z^r + kx^r) \cdot (-x^q) = 0. \quad (6)$$

We next analyze the possibility that we have a singular point when  $xyz = 0$ .

If  $x = 0$ , then equation (4) yields  $z = 0$ , which is then employed in (6) to derive  $y = 0$ , contradiction.

If  $y = 0$ , then equation (5) yields  $x = 0$  and then equation (4) yields  $z = 0$ , contradiction.

If  $z = 0$ , then equation (5) yields  $x = 0$  and then equation (6) yields  $y = 0$ , contradiction.

So, the only possible singular points are of the form  $[x : 1 : z]$ .

We search for possible singular points  $[x : 1 : z] \in \mathbb{P}^2(\mathbb{F}_q)$ . Then equations (4), (5) and (6) read:

$$-x^r + z^{r+1} + kx^r z = 0 \quad (7)$$

$$x^{r+1} - z = 0 \quad (8)$$

$$1 - z^r x - kx^{r+1} = 0. \quad (9)$$

Substituting  $z = x^{r+1}$  from equation (8) into equations (7) and (9), we obtain

$$-x^r + x^{r^2+2r+1} + kx^{2r+1} = 0 \text{ and } 1 - x^{r^2+r+1} - kx^{r+1} = 0,$$

that is, there exists a singular  $\mathbb{F}_q$ -rational point on  $C_{k,r}$  if and only if there exists  $x \in \mathbb{F}_q^*$  such that

$$x^{r^2+r+1} + kx^{r+1} - 1 = 0, \quad (10)$$

as desired.

We end the proof by mentioning that some care is needed to treat the case when the characteristic  $p$  of the field divides the degree of the curve (i.e.,  $p$  divides  $r+1$  in this setting). Indeed, the singular locus of any projective curve  $\{f = 0\}$  is defined by  $\{f = \frac{\partial f}{\partial x} = \frac{\partial f}{\partial y} = \frac{\partial f}{\partial z} = 0\}$ . When  $p$  divides  $\deg(f)$ , it is *not* enough to consider the points in the locus  $\{\frac{\partial f}{\partial x} = \frac{\partial f}{\partial y} = \frac{\partial f}{\partial z} = 0\}$ . Fortunately, in our case, the  $\mathbb{F}_q$ -point  $[x : 1 : z]$  is automatically on the curve  $C_{k,r}$  because  $C_{k,r}$  is plane-filling.  $\square$

It may be natural to make a prediction identical to Conjecture 1.5 for higher-degree curves. However, some care is needed, as the following two examples show. We found these examples using Macaulay2 [GS].

**Example 4.1.** Let  $r = 5$ ,  $q = 11$ , and  $k = 9$ . The plane-filling curve  $C_{9,5}$  over  $\mathbb{F}_{11}$  is smooth at all the  $\mathbb{F}_{11}$ -points because the polynomial  $x^{31} + 9x^6 - 1$  is an irreducible polynomial over  $\mathbb{F}_{11}$ . However,  $C_{9,5}$  is singular at two Galois-conjugate  $\mathbb{F}_{11^2}$ -points.

In the previous example, the curve  $C_{9,5}$  is irreducible over  $\mathbb{F}_{11}$ . Thus,  $C_{9,5}$  satisfies the two conditions of Theorem 3.1 and yet it is singular at two  $\mathbb{F}_{11^2}$ -points. Since  $\deg(C_{9,5}) = q + 6$ , we see that Remark 3.2 is close to being sharp.

**Example 4.2.** Let  $r = 7$ ,  $q = 5$ . In this case, the plane-filling curve  $C_{k,7}$  defined over  $\mathbb{F}_5$  is singular for each  $k \in \mathbb{F}_5$ . Indeed, the associated polynomial  $x^{5^7} + kx^8 - 1$  has an  $\mathbb{F}_5$ -root for  $k \in \{0, 2, 3, 4\}$ . For these values of  $k$ , the curve  $C_{k,r}$  is singular at an  $\mathbb{F}_5$ -point. For  $k = 1$ , the curve  $C_{1,7}$  is singular at four points, namely, two pairs of Galois-conjugate  $\mathbb{F}_{5^2}$ -points.

The two examples above illustrate that Conjecture 1.5 needs to be modified for plane-filling curves of degree  $q+r+1$  when  $r$  is arbitrary. We propose two related conjectures on the smoothness of the curve  $C_{k,r}$  from Theorem 1.8. Recall that  $C_{k,r} \subset \mathbb{P}^2$  is defined by

$$x^r(x^q y - x y^q) + y^r(y^q z - y z^q) + (z^r + kx^r)(z^q x - z x^q) = 0$$

where  $r \geq 2$  is a positive integer and  $k \in \mathbb{F}_q$ .

**Conjecture 4.3.** *Let  $r \geq 2$ . There exists an integer  $m := m(r)$  with the following property. For all finite fields  $\mathbb{F}_q$  with cardinality  $q > m$  and characteristic not dividing  $r$ , there exists some  $k \in \mathbb{F}_q$  such that the curve  $C_{k,r}$  is smooth.*

Using Macaulay2 [GS], we enumerated through values of  $r$  in the range  $[2, 17]$  and  $q$  in the range  $[2, 100]$  with  $\gcd(r, q) = 1$ . We found only the following pairs  $(r, q)$  for which  $C_{k,r}$  is singular for every  $k \in \mathbb{F}_q$ :  $(r, q) = (7, 5), (13, 3), (16, 9),$  and  $(17, 7)$ .

**Conjecture 4.4.** *Let  $r \geq 2$ . There exists an integer  $s := s(r)$  with the following property. For all finite fields  $\mathbb{F}_q$  with characteristic not dividing  $r$ , and for all  $k \in \mathbb{F}_q$ , if  $C_{k,r}$  is smooth at all of its  $\mathbb{F}_{q^s}$ -points, then  $C_{k,r}$  is smooth.*

As a motivation for Conjecture 4.4, we mention the following general fact about pencils of plane curves. The family of plane curves  $C_k$  forms a *pencil* of plane curves since the parameter  $k \in \mathbb{F}_q$  appears linearly in the defining equation. If  $\mathcal{L}$  is a pencil of plane curves in  $\mathbb{P}^2$  parametrized by  $\mathbb{A}^1$ , then  $\mathbb{F}_q$ -members of  $\mathcal{L}$  are defined by  $f(x, y, z) + kg(x, y, z) = 0$  where  $k \in \mathbb{F}_q$  is arbitrary. We will use  $X_k$  to denote this plane curve in the following proposition.

**Proposition 4.5.** *Let  $\mathcal{L}$  be a pencil of plane curves  $\{X_k\}_{k \in \mathbb{F}_q}$  of degree  $d$  defined over a finite field  $\mathbb{F}_q$ . Suppose that for every  $s \geq 1$ , there exists some  $k \in \mathbb{F}_q$  such that  $X_k$  is smooth at all of its  $\mathbb{F}_{q^s}$ -points. Then there exists some  $\ell \in \mathbb{F}_q$  such that  $X_\ell$  is smooth.*

*Proof.* Assume, to the contrary, that  $X_k$  is singular for each  $k \in \mathbb{F}_q$ . For each  $k \in \mathbb{F}_q$ , let  $n_k \in \mathbb{N}$  such that the curve  $X_k$  is singular at some  $\mathbb{F}_{q^{n_k}}$ -point. Let  $N := \prod_{k \in \mathbb{F}_q} n_k$ . By construction, no  $X_k$  is smooth at all of its  $\mathbb{F}_{q^N}$ -points, contradicting the hypothesis.  $\square$

Proposition 4.5 asserts that to find a smooth member of any pencil  $\mathcal{L}$  defined over  $\mathbb{F}_q$ , it is sufficient to find a member which is smooth at all points of an (arbitrary) finite degree. Conjecture 4.4 strengthens the conclusion by predicting that for a pencil of plane-filling curves, one finds a smooth member by only checking smoothness at all points of *fixed* finite degree.

## REFERENCES

- [AG23] Shamil Asgarli and Dragos Ghioca, *Tangent-filling plane curves over finite fields*, Bull. Aust. Math. Soc. (2023), published online on May 2, 2023, available at <https://arxiv.org/abs/2302.13420>.
- [AGY23] Shamil Asgarli, Dragos Ghioca, and Chi Hoi Yip, *Plane curves giving rise to blocking sets over finite fields*, Designs, Codes and Cryptography (2023), to appear, available at <https://arxiv.org/abs/2208.13299>.
- [AP96] Yves Aubry and Marc Perret, *A Weil theorem for singular curves*, Arithmetic, geometry and coding theory (Luminy, 1993), 1996, pp. 1–7.
- [DC18] Gregory Duran Cunha, *Curves containing all points of a finite projective Galois plane*, J. Pure Appl. Algebra **222** (2018), no. 10, 2964–2974.
- [Gab01] O. Gabber, *On space filling curves and Albanese varieties*, Geom. Funct. Anal. **11** (2001), no. 6, 1192–1200.
- [Hom20] Masaaki Homma, *Fragments of plane filling curves of degree  $q + 2$  over the finite field of  $q$  elements, and of affine-plane filling curves of degree  $q + 1$* , Linear Algebra Appl. **589** (2020), 9–27.
- [HK13] Masaaki Homma and Seon Jeong Kim, *Nonsingular plane filling curves of minimum degree over a finite field and their automorphism groups: supplements to a work of Tallini*, Linear Algebra Appl. **438** (2013), no. 3, 969–985.
- [HK23] ———, *Filling curves for  $\mathbb{P}^1 \times \mathbb{P}^1$* , Comm. Algebra **51** (2023), no. 6, 2680–2687.
- [Kat99] Nicholas M. Katz, *Space filling curves over finite fields*, Math. Res. Lett. **6** (1999), no. 5-6, 613–624.
- [GS] Daniel R. Grayson and Michael E. Stillman, *Macaulay2, a software system for research in algebraic geometry*. <http://www.math.uiuc.edu/Macaulay2/>.
- [Pea90] Giuseppe Peano, *Sur une courbe, qui remplit toute une aire plane*, Math. Ann. **36** (1890), no. 1, 157–160.
- [Poo04] Bjorn Poonen, *Bertini theorems over finite fields*, Ann. of Math. (2) **160** (2004), no. 3, 1099–1127.
- [Tal61a] Giuseppe Tallini, *Le ipersuperficie irriducibili d'ordine minimo che invadono uno spazio di Galois*, Atti Accad. Naz. Lincei Rend. Cl. Sci. Fis. Mat. Nat. (8) **30** (1961), 706–712.
- [Tal61b] ———, *Sulle ipersuperficie irriducibili d'ordine minimo che contengono tutti i punti di uno spazio di Galois  $S_{r,q}$* , Rend. Mat. e Appl. (5) **20** (1961), 431–479.

DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCE, SANTA CLARA UNIVERSITY, 500 EL CAMINO REAL, USA 95053

*Email address:* sasgarli@scu.edu

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF BRITISH COLUMBIA, 1984 MATHEMATICS ROAD, CANADA V6T 1Z2

*Email address:* dghioca@math.ubc.ca