

## Math 223: Problem Set 4 (due 3/10/12)

### Practice problems (recommended, but do not submit)

Section 2.1, Problems 1-3,5,9,10-12,28-29

Section 2.2, Problems 1-3.

### Calculations with linear maps

1. Let  $T: U \rightarrow V$  be a linear map, and let  $S \subset U$  be a generating set. Show that  $\{T\underline{s} \mid \underline{s} \in S\}$  is a generating set for  $\text{Im } T$ .

RMK This is a starting point for finding a basis for  $\text{Im } T$ .

2. Let  $T: \mathbb{R}^2 \rightarrow \mathbb{R}^3$  be the linear map  $T \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} x_1 + x_2 \\ x_1 - x_2 \\ 2x_1 \end{pmatrix}$ .

(a) Find bases for  $\text{Ker } T$ ,  $\text{Im } T$  and check that the dimension formula holds.

(b) Find the matrix for  $T$  with respect to the bases  $\left\{ \begin{pmatrix} 1 \\ 1 \end{pmatrix}, \begin{pmatrix} 1 \\ -1 \end{pmatrix} \right\}$  of  $\mathbb{R}^2$  and  $\left\{ \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix}, \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix}, \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} \right\}$  of  $\mathbb{R}^3$ .

3. Let  $T: \mathbb{R}^5 \rightarrow \mathbb{R}^3$  be the linear map  $T \begin{pmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \\ x_5 \end{pmatrix} = \begin{pmatrix} 2x_1 + x_2 \\ x_1 - x_2 + x_3 - x_5 \\ -3x_1 - x_3 + x_5 \end{pmatrix}$ .

(a) Find bases for  $\text{Ker } T$ ,  $\text{Im } T$  (use problem 1) and check that the dimension formula holds.

(b) Find the matrix for  $T$  with respect to the standard bases of  $\mathbb{R}^5$ ,  $\mathbb{R}^3$ .

(c) Find the matrix for  $T$  with respect to the standard basis of  $\mathbb{R}^5$  and the basis  $\left\{ \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix}, \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix} \right\}$  of  $\mathbb{R}^3$ .

4. Let  $D: \mathbb{R}[x]^{\leq n} \rightarrow \mathbb{R}[x]^{\leq n}$  be the differentiation map.

(a) Find  $\text{Ker } D$  and its dimension.

(b) Find  $\text{Im } D$ .

Fix a number  $a \neq 0$  and let  $T: \mathbb{R}[x]^{\leq n} \rightarrow \mathbb{R}[x]^{\leq n}$  be the map  $D + Za$  (that is,  $Tp = \frac{dp}{dx} + a \cdot p$ ).

(c) Show that  $T$  maps the basis of monomials to a set of  $n+1$  polynomials of distinct degrees.

(\*d) Show that  $\text{Im } T = \mathbb{R}[x]^{\leq n}$ .

### Linear dependence of functions

5. Let  $X$  be a set, and let  $\{f_i\}_{i=1}^n \subset \mathbb{R}^X$  be some  $n$  functions. Let  $\{x_j\}_{j=1}^m \subset X$  be  $m$  points of  $X$ .

(a) Define a map  $E: \mathbb{R}^n \rightarrow \mathbb{R}^m$  by setting  $(E\underline{a})_j = \sum_{i=1}^n a_i f_i(x_j)$  for  $\underline{a} \in \mathbb{R}^n$  and  $1 \leq j \leq m$ . Show that  $E$  is linear.

(b) Suppose that  $m < n$ . Show that  $\dim \text{Ker } E > 0$ . Conclude that if  $m < n$  there exist  $\{a_i\}_{i=1}^n$  not all zero such that the function  $\sum_{i=1}^n a_i f_i$  vanishes at all the points  $\{x_j\}_{j=1}^m$ .

### Surjective and injective maps; Invertibility

DEFINITION. Let  $T: U \rightarrow V$  be a linear map. We say that  $T$  is *injective* (a *monomorphism*) if  $T\underline{u} = T\underline{u}'$  implies  $\underline{u} = \underline{u}'$  and *surjective* (an *epimorphism*) if  $\text{Im} T = V$ .

6. Show that  $T$  is injective if and only if  $\text{Ker} T = \{\underline{0}\}$ . (Hint: to compare two vectors consider their difference)

DEFINITION. If a linear map  $T: U \rightarrow V$  is surjective and injective we say it is an *isomorphism* (of vector spaces). We say that  $U, V$  are isomorphic if there is an isomorphism between them.

7. Suppose that  $T: U \rightarrow V$  is an isomorphism of vector spaces, and define a function  $T^{-1}: V \rightarrow U$  by  $T^{-1}\underline{v}$  being that vector  $\underline{u}$  such that  $T\underline{u} = \underline{v}$ .
- (a) Explain why  $\underline{u}$  exists and why it is unique (that is, review the definitions of surjective and injective)
- (\*b) Show that  $T^{-1}$  is a linear function.

---

### Supplementary problems

- A. Let  $V$  be a vector space and let  $W_1, W_2 \subset V$  be finite-dimensional subspaces.
- (a) Show that  $\dim(W_1 + W_2) \leq \dim W_1 + \dim W_2$ .
- (\*\*b) Show that  $\dim(W_1 + W_2) + \dim(W_1 \cap W_2) = \dim W_1 + \dim W_2$ .
- RMK Let  $A, B$  be finite sets. Then the “inclusion-exclusion” formula states  $\#A + \#B = \#(A \cup B) + \#(A \cap B)$
- B. Let  $V$  be a vector space,  $W$  a subspace. Let  $B \subset W$  be a basis for  $W$  and let  $C \subset V$  be such that  $B \cup C$  is a basis for  $V$  (that is, we extend  $B$  until we get a basis for  $V$ ).
- (a) Show that  $\{\underline{v} + W\}_{\underline{v} \in C}$  is a basis for the quotient vector space  $V/W$  (see supplement to PS2).
- (b) Conclude that  $\dim V = \dim V/W + \dim W$ .
- (c) Show that the map  $v \mapsto v + W$  gives a surjective linear map  $V \rightarrow V/W$  with kernel  $W$ .