

## Math 322: Problem Set 6 (due 27/10/2015)

### Practice problems

- P1. Let  $G$  be a group and let  $X$  be a set of size at least 2. Fix  $x_0 \in X$  and for  $g \in G, x \in X$  set  $g \cdot x = x_0$ .
- Show that this operation satisfies  $(gh) \cdot x = g \cdot (h \cdot x)$  for all  $g, h \in G, x \in X$ .
  - This is not a group action. Why?
- P2. Let  $G$  act on  $X$ . Say that  $A \subset X$  is  $G$ -invariant if for every  $g \in G, a \in A$  we have  $g \cdot a \in A$ .
- Show that  $A$  is  $G$ -invariant iff  $g \cdot A = A$  ( $g \cdot A$  in the sense of problem 4(a)).
  - Suppose  $A$  is  $G$ -invariant. Show that the restriction of the action to  $A$  (formally, the binary operation  $\cdot \upharpoonright_{G \times A}$ ) is an action of  $G$  on  $A$ .

### Simplicity of $A_n$

- Let  $V = \{\text{id}, (12)(34), (13)(24), (14)(23)\}$ . Show that  $V \triangleleft S_4$ , so that  $S_4$  is not simple.
- (The normal subgroups of  $S_n$ ) Let  $N \triangleleft S_n$  with  $n \geq 5$ .
  - Let  $G$  be a group and let  $H \triangleleft G$  be a normal subgroup isomorphic to  $C_2$ . Show that  $H < Z(G)$ .
  - Suppose that  $N \cap A_n \neq \{\text{id}\}$ . Show that  $N \supset A_n$  and conclude that  $N = A_n$  or  $N = S_n$ .
  - Suppose that  $N \cap A_n = \{\text{id}\}$ . Show that  $N$  is isomorphic to a subgroup of  $C_2$ .
  - Show that if  $n \geq 3$  then  $Z(S_n) = \{\text{id}\}$ , and conclude that in case (c) we must have  $N = \text{id}$ .
- Let  $X$  be an infinite set.
  - Show that  $S_X^{\text{fin}} = \{\sigma \in S_X \mid \text{supp}(\sigma) \text{ is finite}\}$  is a subgroup of  $S_X$ .  
PRAC For finite  $F \subset X$  there is a natural inclusion  $S_F \hookrightarrow S_X$ , which is a group homomorphism and an isomorphism onto its image. Let  $\text{sgn}_F: S_F \rightarrow \{\pm 1\}$  be the sign character.  
DEF For  $\sigma \in S_X^{\text{fin}}$  define  $\text{sgn}(\sigma) = \text{sgn}_F(\sigma)$  for any finite  $F$  such that  $\sigma \in S_F$ .
  - Show that  $\text{sgn}(\sigma)$  is well-defined (independent of  $F$ ) and a group hom  $S_X^{\text{fin}} \rightarrow \{\pm 1\}$ .
  - (\*d) The *infinite alternating group*  $A_X$  is kernel of this homomorphism. Show that  $A_X$  is simple.

### Group actions

- Let the group  $G$  act on the set  $X$ .
  - For  $g \in G$  and  $A \in P(X)$  set  $g \cdot A = \{g \cdot a \mid a \in A\} = \{x \in X \mid \exists a \in A : x = g \cdot a\}$ . Show that this defines an action of  $G$  on  $P(X)$ .
  - In PS2 we endowed  $P(X)$  with a group structure. Show that the action of (a) is by *automorphisms*: that the map  $A \mapsto g \cdot A$  is a group homomorphism  $(P(X), \Delta) \rightarrow (P(X), \Delta)$ .
  - Let  $Y$  be another set. For  $f: X \rightarrow Y$  set  $(g \cdot f)(x) = f(g^{-1} \cdot x)$ . Show that this defines an action of  $G$  on  $Y^X$ , the set of functions from  $X$  to  $Y$ .
  - (\*d) Suppose that  $Y = \mathbb{R}$  (or any field), so that  $\mathbb{R}^X$  has the structure of a vector space over  $\mathbb{R}$ . Show that the action of (c) is by *linear maps*.

5. (Some stabilizers) The action of  $S_X$  on  $X$  induces an action on  $P(X)$  as in problem 4(a). Suppose that  $X$  is finite,  $\#X = n$ .
- (a) Show that the orbits of  $S_X$  on  $P(X)$  are the sets  $\binom{X}{k} = \{A \subset X \mid \#A = k\}$  for  $0 \leq k \leq n$ .  
 SUPP When  $X$  is infinite,  $\binom{X}{\kappa}$  are orbits if  $\kappa < |X|$ , but there are multiple orbits on  $\binom{X}{|X|}$ , parametrized by the cardinality of the complement.
- (b) Let  $A \subset X$ . Show that  $\text{Stab}_{S_X}(A) \simeq S_A \times S_{X-A}$ .
- (c) Use (a),(b) to show that  $\#\binom{X}{k} = \frac{n!}{k!(n-k)!}$ .

### Supplementary problem: Conjugation

- A. Let  $G$  be a finite groups.
- (a) Suppose all elements of  $G$  are conjugate. Show that  $G = \{e\}$ .
- (b) Suppose  $G$  has exactly two conjugacy classes. Show that  $G \simeq C_2$ .
- (\*c) Suppose  $G$  has exactly three conjugacy classes. Show that  $G \simeq C_3$ .
- (\*\*d) Show that for each  $k$  there is  $N = N(k)$  such that if  $G$  has at most  $k$  conjugacy classes its order is at most  $N$ .

RMK There exists an infinite group in which all non-identity elements are conjugate.

(hint for 2(a): let  $H = \{1, h\}$ , let  $g \in G$ , and consider the element  $ghg^{-1}$ )

(hint for 2(b): consider the index of  $N$ )

(hint for 2(c): restrict  $\text{sgn}: S_n \rightarrow C_2$  to  $N$ )

(hint for 6(b): the number of conjugates of an element divides the order of the group)