

Math 101 – SOLUTIONS TO WORKSHEET 17
IMPROPER INTEGRALS

1. IMPROPER AT INFINITY

- (1) For which values of p does $\int_1^\infty \frac{1}{x^p} dx$ converge? Diverge?

Solution: For $p \neq -1$ we have $\int_1^T \frac{1}{x^p} dx = \left[\frac{1}{1-p} x^{1-p} \right]_1^T = \frac{1}{1-p} (T^{1-p} - 1)$. If $1 - p > 0$ then $T^{1-p} \rightarrow \infty$ so the integral diverges. If $1 - p < 0$ then $T^{1-p} \rightarrow 0$ and the integral converges. If $1 - p = 0$ then $\int_1^T \frac{1}{x^p} dx = [\log x]_1^T = \log T \rightarrow \infty$ and the integral diverges.

- (2) (Final, 2010) Evaluate $\int_{-\infty}^{-1} e^{2x} dx$. Simplify your answer as much as possible.

Solution: $\int_T^{-1} e^{2x} dx = \left[\frac{1}{2} e^{2x} \right]_{x=T}^{x=-1} = \frac{1}{2} (e^{-2} - e^{2T})$. Now $\lim_{T \rightarrow -\infty} e^{2T} = \lim_{x \rightarrow -\infty} e^x = 0$ so

$$\int_{-\infty}^{-1} e^{2x} dx = \lim_{T \rightarrow -\infty} \frac{1}{2} (e^{-2} - e^{2T}) = \boxed{\frac{1}{2} e^{-2}}.$$

Solution: $\int_{-T}^{-1} e^{2x} dx = \left[\frac{1}{2} e^{2x} \right]_{x=-T}^{x=-1} = \frac{1}{2} (e^{-2} - e^{-2T})$. Now $\lim_{T \rightarrow \infty} e^{-2T} = \lim_{x \rightarrow \infty} e^{-x} = 0$ so

$$\int_{-\infty}^{-1} e^{2x} dx = \lim_{T \rightarrow \infty} \frac{1}{2} (e^{-2} - e^{-2T}) = \boxed{\frac{1}{2} e^{-2}}.$$

- (3) Find a constant C such that $\int_{-\infty}^{+\infty} \frac{C dx}{1+x^2} = 1$.

Solution: $\int_0^T \frac{C dx}{1+x^2} = C [\arctan x]_{x=0}^{x=T} = C \arctan T$ so $\int_0^\infty \frac{dx}{1+x^2} = C \lim_{T \rightarrow \infty} \arctan T = \frac{\pi}{2} C$. Since the function is symmetric we also have $\int_{-\infty}^0 \frac{C dx}{1+x^2} = \frac{\pi}{2} C$ so

$$\int_{-\infty}^{+\infty} \frac{C dx}{1+x^2} = \pi C$$

and we need to take $C = \frac{1}{\pi}$.

- (4) We study $\int_{-\infty}^{+\infty} x dx$.

(a) Evaluate $\int_{-T}^T x dx$.

(b) Evaluate $\lim_{T \rightarrow \infty} \int_{-T}^T x dx$.

(c) Does the integral converge?

Solution: $\int_T^{-T} x dx = 0$ since the integrand is odd, so $\lim_{T \rightarrow \infty} \int_{-T}^T x dx = 0$. Nevertheless the integral diverges: $\int_0^\infty x dx$ diverges, as does $\int_{-\infty}^0 x dx$.

- (5) (Final, 2009) For what values of p does $\int_e^\infty \frac{dx}{x(\log x)^p}$ converge?

Solution: We note that $\frac{dx}{x} = d(\log x)$, so letting $u = \log x$ we have $\int_{x=e}^{x=T} \frac{dx}{x(\log x)^p} = \int_1^{\log T} \frac{du}{u^p}$. Now as $T \rightarrow \infty$, $\log T \rightarrow \infty$ as well, so this converges exactly when $\int_1^\infty \frac{du}{u^p}$ converges, that is exactly for $\boxed{p > 1}$.

2. IMPROPER AT FINITE POINTS

- (6) For which values of p does $\int_0^1 \frac{dx}{x^p}$ converge?

Solution: For $p \neq 1$ we have $\int_\epsilon^1 \frac{dx}{x^p} = \left[\frac{x^{1-p}}{1-p} \right]_\epsilon^1 = \frac{1}{1-p} - \frac{\epsilon^{1-p}}{1-p}$. Now as $\epsilon \rightarrow 0$,

$$\lim_{\epsilon \rightarrow 0} \epsilon^{1-p} = \begin{cases} 0 & 1-p > 0 \\ \infty & 1-p < 0 \end{cases}$$

so $\int_0^1 \frac{dx}{x^p}$ exists when $p < 1$ and diverges when $p > 1$. For $p = 1$ we have $\int_\epsilon^1 \frac{dx}{x} = -\log \epsilon \xrightarrow{\epsilon \rightarrow 0} \infty$ and the integral diverges as well.

- (7) (Math 103 Final, 2013) Evaluate the integral if it exists, otherwise show that it doesn't: $I = \int_0^2 \frac{dx}{1-x^2}$.

Solution: The function $\frac{1}{1-x^2}$ is discontinuous at $x = 1$, so we need to consider the convergence of $\int_0^1 \frac{dx}{1-x^2}$ and $\int_1^2 \frac{dx}{1-x^2}$ separately. Considering the first integral, for $0 \leq x < 1$ we have $\frac{1}{1-x^2} = \frac{1}{(1-x)(1+x)} \geq \frac{1}{1-x} \cdot \frac{1}{1+1}$. Now letting $1-x = u$ we have $\int_{x=0}^{x=1} \frac{dx}{2(1-x)} = \frac{1}{2} \int_{u=0}^{u=1} \frac{du}{u}$ diverges by the p -test ($p = 1$) so our integral diverges.

Solution: We have $\frac{1}{1-x^2} = \frac{1}{2} \frac{1}{1-x} + \frac{1}{2} \frac{1}{1+x}$. The second function is continuous on $[0, 2]$ so it's enough to study $\frac{1}{2} \int_0^2 \frac{dx}{1-x}$. Changing variables to $u = 1-x$ we have

$$\int_{x=0}^{x=2} \frac{dx}{1-x} = \int_{u=1}^{u=-1} \frac{-du}{u} = \int_{u=-1}^{u=1} \frac{du}{u}.$$

Now both $\int_{-1}^0 \frac{du}{u}$ and $\int_0^1 \frac{du}{u}$ diverge by the p -test so the integral diverges.

3. COMPARISON OF INTEGRALS

- (8) Decide which of the following integrals converge

- (a) (103 Final, 2012) $\int_1^\infty \frac{1+\sin x}{x^2} dx$.

Solution: Let $g(x) = \frac{1+\sin x}{x^2}$. Since $\sin x \geq -1$ for all x we have $g(x) \geq \frac{1+(-1)}{x^2} = 0$. Since $\sin x \leq 1$ for all x we have $g(x) \leq \frac{1+1}{x^2} = \frac{2}{x^2}$. Now $\int_1^\infty \frac{dx}{x^2}$ converges by the p -test ($p = 2 > 1$) so $\int_1^\infty \frac{2dx}{x^2}$ converges as well. By the comparison test it follows that $\int_1^\infty g(x) dx$ converges.

- (b) $\int_1^\infty \frac{3-\cos x}{x} dx$.

Solution: Let $f(x) = \frac{3+\cos x}{x}$. $\cos x \geq -1$ we for all x we have $f(x) \geq \frac{3+(-1)}{x} \geq \frac{1}{x}$. Now $\int_1^\infty \frac{dx}{x}$ diverges by the p -test ($p = 1 > 1$) so by the comparison test $\int_1^\infty \frac{3+\cos x}{x} dx$ diverges as well.

- (c) (Bell curve) $\int_{-\infty}^{+\infty} e^{-x^2} dx$

Solution: By symmetry (the function is even) it's enough to consider $\int_0^\infty e^{-x^2} dx$. Since $\int_0^1 e^{-x^2} dx$ exists, it's enough to consider $\int_1^\infty e^{-x^2} dx$. But for $x \geq 1$, $x^2 \geq x$ so $e^{-x^2} \leq e^{-x}$. Now $\int_1^\infty e^{-x} dx$ converges (exponential function fact) and $e^{-x^2} \geq 0$ for all x , so by the comparison test $\int_1^\infty e^{-x^2} dx$ converges. It follows that $\int_{-\infty}^{+\infty} e^{-x^2} dx$ converges as well.

- (d) $\int_0^1 \frac{dx}{\sqrt{x+\sin x}}$

Solution: The function is continuous on $(0, 1]$ and since $1 < \pi$ we have $0 < \sin x$ on that interval. It follows that $\sqrt{x} + \sin x > \sqrt{x} > 0$ and thus $0 < \frac{1}{\sqrt{x+\sin x}} < \frac{1}{\sqrt{x}}$ on our interval. Now $\int_0^1 \frac{dx}{\sqrt{x}}$ converges by the p -test ($p = \frac{1}{2} < 1$) so by the comparison test the given integral converges as well.

- (e) (hard) $\int_0^1 \frac{dx}{x^2+x^3}$

Solution: Multiplying the inequality $0 < x \leq 1$ by x^2 we see that $x^3 < x^2$. It follows that $x^2 + x^3 \leq 2x^2$ so that $\frac{1}{x^2+x^3} \geq \frac{1}{2x^2}$. Now $\int_0^1 \frac{dx}{2x^2} = \frac{1}{2} \int_0^1 \frac{dx}{x^2}$ diverges by the p -test ($p = 2 > 1$) so by the comparison test the given integral diverges as well.

- (f) (hard) $\int_0^\infty \frac{x^{1000}}{e^x} dx$

Solution: Any exponential function grows faster than any polynomial function, so that $x^{1000} \leq e^{x/2}$ for x large enough (1000 applications of l'Hôpital's rule will show $\lim_{x \rightarrow \infty} \frac{x^{1000}}{e^{x/2}} = 0$). It follows that $\frac{x^{1000}}{e^x} = \frac{x^{1000}}{e^{x/2}} \cdot \frac{1}{e^{x/2}} \leq \frac{1}{e^{x/2}}$ for x large enough. But $\int_0^\infty \frac{1}{e^{x/2}} dx$ exists (exponential decay), so the given integral exists as well.