

Lior Silberman's Math 412: Problem Set 5 (due 12/10/2017)

Practice

- P1. Let $U = \text{Span}_F \{u_1, u_2\}$ be two-dimensional. Show that the element $\underline{u}_1 \otimes \underline{u}_1 + \underline{u}_2 \otimes \underline{u}_2 \in U \otimes U$ is not a pure tensor, that is not of the form $\underline{u} \otimes \underline{u}$ for any $\underline{u} \in U$.
- P2. Let $\iota: U \times V \rightarrow U \otimes V$ be the standard inclusion map ($\iota(\underline{u}, \underline{v}) = \underline{u} \otimes \underline{v}$). Show that $\iota(\underline{u}, \underline{v}) = 0$ iff $\underline{u} = \underline{0}_U$ or $\underline{v} = \underline{0}_V$ and that for non-zero vectors we have $\iota(\underline{u}, \underline{v}) = \iota(\underline{u}', \underline{v}')$ iff $\underline{u}' = \alpha \underline{u}$ and $\underline{v}' = \alpha^{-1} \underline{v}$ for some $\alpha \in F^\times$.
- P3. Let U, V be finite-dimensional spaces and let $A \in \text{End}(U), B \in \text{End}(V)$.
 (a) Construct a map $A \oplus B \in \text{End}_F(U \oplus V)$ restricting to A, B on the images of U, V in $U \oplus V$.
 (b) Show that $\text{Tr}(A \oplus B) = \text{Tr}(A) + \text{Tr}(B)$.
 (c) Evaluate $\det(A \oplus B)$.

Tensor products of maps

1. Let U, V be finite-dimensional spaces, and let $A \in \text{End}(U), B \in \text{End}(V)$.
 (a) Show that $(\underline{u}, \underline{v}) \mapsto (A\underline{u}) \otimes (B\underline{v})$ is bilinear, and obtain a linear map $A \otimes B \in \text{End}(U \otimes V)$.
 (b) Suppose A, B are diagonalizable. Using an appropriate basis for $U \otimes V$, Obtain a formula for $\det(A \otimes B)$ in terms of $\det(A)$ and $\det(B)$.
 (c) Extending (a) by induction, show for any $A \in \text{End}_F(V)$, the map $A^{\otimes k}$ induces maps $\text{Sym}^k A \in \text{End}(\text{Sym}^k V)$ and $\wedge^k A \in \text{End}(\wedge^k V)$.
 (*d) Show that the formula of (b) holds for all A, B .
 SUPP (Notation continued from supplement to PS4) Let $V_K = K \otimes_F V$ be an extension of scalars. For $T \in \text{End}_F(V)$ let $T_K = \text{Id}_K \otimes T$. Show that $T_K \in \text{End}_K(V_K)$, and that the natural inclusions $\text{Ker}(T), \text{Im}(T) \subset V$ extend to identifications $(\text{Ker}(T))_K = \text{Ker}(T_K)$ and $(\text{Im}(T))_K = \text{Im}(T_K)$.

2. Suppose $\frac{1}{2} \in F$, and let U be finite-dimensional. Construct isomorphisms

$$\{\text{symmetric bilinear forms on } U\} \leftrightarrow (\text{Sym}^2 U)' \leftrightarrow \text{Sym}^2(U').$$

Nilpotence

3. Let $U \in M_n(F)$ be *strictly upper-triangular*, that is upper triangular with zeroes along the diagonal. Show that $U^n = 0$ and construct such U with $U^{n-1} \neq 0$.
4. Let V be a finite-dimensional vector space, $T \in \text{End}(V)$.
 (*a) Show that the following statements are equivalent:
 (1) $\forall \underline{v} \in V : \exists k \geq 0 : T^k \underline{v} = \underline{0}$; (2) $\exists k \geq 0 : \forall \underline{v} \in V : T^k \underline{v} = \underline{0}$.
 DEF A linear map satisfying (2) is called *nilpotent*. Example: see problem 5.
 (b) Find nilpotent $A, B \in M_2(F)$ such that $A + B$ isn't nilpotent.
 (c) Suppose that $A, B \in \text{End}(V)$ are nilpotent and that A, B commute. Show that $A + B$ is nilpotent.

Extra credit

5. Let V be finite-dimensional.
- Construct an isomorphism $U \otimes V' \rightarrow \text{Hom}_F(V, U)$.
 - Define a map $\text{Tr}: U \otimes U' \rightarrow F$ extending the evaluation pairing $U \times U' \rightarrow F$.
- DEF The *trace* of $T \in \text{Hom}_F(U, U)$ is given by identifying T with an element of $U \otimes U'$ via (a) and then applying the map of (b).
- Let $T \in \text{End}_F(U)$, and let A be the matrix of T with respect to the basis $\{u_i\}_{i=1}^n \subset U$. Show that $\text{Tr} T = \sum_{i=1}^n A_{ii}$.
- RMK This shows that similar matrices have the same trace!
- Solve P3(b) from this point of view.

Supplementary problems

- A. (The tensor algebra) Fix a vector space U .
- Extend the bilinear map $\otimes: U^{\otimes n} \times U^{\otimes m} \rightarrow U^{\otimes n} \otimes U^{\otimes m} \simeq U^{\otimes(n+m)}$ to a bilinear map $\otimes: \bigoplus_{n=0}^{\infty} U^{\otimes n} \times \bigoplus_{n=0}^{\infty} U^{\otimes n} \rightarrow \bigoplus_{n=0}^{\infty} U^{\otimes n}$.
 - Show that this map \otimes is associative and distributive over addition. Show that $1_F \in F \simeq U^{\otimes 0}$ is an identity for this multiplication.
- DEF This algebra is called the *tensor algebra* $T(U)$.
- Show that the tensor algebra is *free*: for any F -algebra A and any F -linear map $f: U \rightarrow A$ there is a unique F -algebra homomorphism $\tilde{f}: T(U) \rightarrow A$ whose restriction to $U^{\otimes 1}$ is f .
- B. (The symmetric algebra). Fix a vector space U .
- Endow $\bigoplus_{n=0}^{\infty} \text{Sym}^n U$ with a product structure as in 3(a).
 - Show that this creates a commutative algebra $\text{Sym}(U)$.
 - Fixing a basis $\{u_i\}_{i \in I} \subset U$, construct an isomorphism $F[\{x_i\}_{i \in I}] \rightarrow \text{Sym}^* U$.
- RMK In particular, $\text{Sym}^*(U')$ gives a coordinate-free notion of “polynomial function on U ”.
- Let $I \triangleleft T(U)$ be the two-sided ideal generated by all elements of the form $u \otimes v - v \otimes u \in U^{\otimes 2}$. Show that the map $\text{Sym}(U) \rightarrow T(U)/I$ is an isomorphism.
- RMK When the field F has finite characteristic, the correct definition of the symmetric algebra (the definition which gives the universal property) is $\text{Sym}(U) \stackrel{\text{def}}{=} T(U)/I$, not the space of symmetric tensors.
- C. Let V be a (possibly infinite-dimensional) vector space, $A \in \text{End}(V)$.
- Show that the following are equivalent for $v \in V$:
 - $\dim_F \text{Span}_F \{A^n v\}_{n=0}^{\infty} < \infty$;
 - there is a finite-dimensional subspace $W \subset V$ such that $AW \subset W$.
- DEF Call such v *locally finite*, and let V_{fin} be the set of locally finite vectors.
- Show that V_{fin} is a subspace of V .
 - Call A *locally nilpotent* if for every $v \in V$ there is $n \geq 0$ such that $A^n v = \underline{0}$ (condition (1) of 5(a)). Find a vector space V and a locally nilpotent map $A \in \text{End}(V)$ which is not nilpotent.
 - (*d) A is called *locally finite* if $V_{\text{fin}} = V$, that is if every vector is contained in a finite-dimensional A -invariant subspace. Find a space V and locally finite linear maps $A, B \in \text{End}(V)$ such that $A + B$ is not locally finite.