Math 100 – WORKSHEET 5 THE DERIVATIVE

1. LINEAR COMBINATIONS; POWER LAWS

(1) If f, g are functions and a, b are numbers then (af + bg)' = af' + bg'(2) $\frac{d}{dx}(x^r) = rx^{r-1}$ (3) $\frac{d}{dx}(e^x) = e^x$.

(1)

(a) Differentiate $f(x) = \frac{5x^3 - 2x + 1}{\sqrt{x}}$.

(b) Let $g(x) = Ax^{5/2} + x^2$. Suppose that g'(4) = 0. What is A?

- (2) Find the *second* derivative of
 - (a) $5e^x$
 - (b) $\sqrt{x} + 5e^x$

(3) The line y = 5x + B is tangent to the curve $y = x^3 + 2x$. What is B?

Date: 19/9/2019, Worksheet by Lior Silberman. This instructional material is excluded from the terms of UBC Policy 81.

Fact.
$$(fg)' = f'g + fg', \left(\frac{f}{g}\right)' = \frac{f'g - fg'}{g^2}$$

(1) Find $\frac{\mathrm{d}}{\mathrm{d}x}(x^a e^x)$.

(2) Suppose that f(1) = 1, g(1) = 2, f'(1) = 3, g'(1) = 4. Find (fg)'(1) and $\left(\frac{f}{g}\right)'(1)$.

(3)
$$f(x) = \frac{x^2 + A}{\sqrt{x}}$$
. $f'(x) =$

(4) Let $f(x) = \frac{x}{\sqrt{x}+A}$. Given that $f'(4) = \frac{3}{16}$, give a quadratic equation for A.