

**Math 100A – SOLUTIONS TO WORKSHEET 7**  
**CURVE SKETCHING**


1. CONVEXITY AND CONCAVITY

- (1) Consider the curve  $y = x^3 - x$ .
- (a) ★ Find the line tangent to the curve at  $x = 1$ .  
**Solution:**  $\frac{dy}{dx} = 3x^2 - 1$  so the derivative at  $x = 1$  is 2. Since  $y(1) = 0$  the line is  $Y = 2(X - 1)$ .
- (b) ★★ Near  $x = 1$ , is the line above or below the curve? Hint: how does the slope of the curve behave to the right and left of the point?  
**Solution:** Since  $x > 0$  the slope is increasing near  $x = 1$ , so to the right the function grows faster than the line, to the right is decreases slower than the line, and the line is below.  
**Solution:** We have  $x^3 - x - 2(x - 1) = (x - 1)(x^2 + x - 2) = (x - 1)^2(x + 2) = 3(x - 1)^2 + (x - 1)^3$  which is positive for  $x$  close enough to 1. In next week's lecture we'll talk more about the representation  $x^3 - x = 2(x - 1) + 3(x - 1)^2 + (x - 1)^3$ .
- (2) For each curve find its domain; where is it concave up or down? Where are the inflection points.
- (a) ★  $y = x \log x - \frac{1}{2}x^2$ .  
**Solution:** This is defined on  $(0, \infty)$ . We have  $y' = \log x - 1 - x$  so  $y'' = \frac{1}{x} - 1$ . Thus  $y'' > 0$  if  $x < 1$ ,  $y'' < 0$  if  $x > 1$ , and the function is concave up on  $(0, 1)$ , concave down on  $(1, \infty)$  and has an inflection point at  $x = 1$ .
- (b) ★  $y = \sqrt[3]{x}$ .  
**Solution:** This is an odd root, which is defined (and continuous) on the entire line. We have  $y' = \frac{1}{3}x^{-2/3}$  which is defined for  $x \neq 0$  (the tangent line at  $x = 0$  is vertical, as we can see by switching to the representation  $x = y^3$ ). We then have  $y'' = -\frac{2}{9}x^{-5/3}$  which is positive when  $x < 0$  and positive when  $x > 0$ , so the function changes concavity at  $x = 0$  and that is an inflection point.

2. CURVE SKETCHING

- (3) ★★ Let  $f(x) = \frac{x^2}{x^2+1}$  for which  $f'(x) = \frac{2x}{(x^2+1)^2}$  and  $f''(x) = \frac{2(1-3x^2)}{(x^2+1)^3}$ .
- (a) What are the domain and intercepts of  $f$ ? What are the asymptotics at  $\pm\infty$ ? Are there any vertical asymptotes? What are the asymptotics there?  
**Solution:** The function is defined for all  $x$  (always have  $x^2 + 1 > 0$ ). We have  $f(0) = 0$  and conversely if  $f(x) = 0$  then  $x^2 = 0$  so  $x = 0$ . As  $x \rightarrow \pm\infty$  we have
- $$\frac{x^2}{x^2+1} \sim \frac{x^2}{x^2} = 1$$
- so we have the horizontal asymptote  $y = 1$  in both ends.
- (b) What are the intervals of increase/decrease? The local and global extrema?  
**Solution:** Since  $\frac{2}{(x^2+1)^2}$  is always positive,  $f'(x) > 0$  when  $x > 0$  and  $f'(x) < 0$  when  $x < 0$ . Thus  $f$  is decreasing on  $(-\infty, 0)$ , increasing on  $(0, \infty)$  and has a local (and global) minimum at  $x = 0$ .
- (c) What are the intervals of concavity? Any inflection points?  
**Solution:** Since  $\frac{2}{(x^2+1)^3}$  is always positive, the sign of  $f''(x)$  is the same as that of  $1 - 3x^2$ . In particular  $f''(x) > 0$  when  $1 - 3x^2 > 0$ , that is when  $3x^2 < 1$  so when  $|x| < \frac{1}{\sqrt{3}}$ . Conversely  $f''(x) < 0$  when  $1 - 3x^2 < 0$  that is when  $|x| > \frac{1}{\sqrt{3}}$  or when  $x \in \left(-\infty, -\frac{1}{\sqrt{3}}\right) \cup \left(\frac{1}{\sqrt{3}}, \infty\right)$ . We thus have inflection points at  $\pm\frac{1}{\sqrt{3}}$ .

(d) Sketch a graph of  $f(x)$ .



(4) **★★** Let  $f(x) = \frac{1}{\sqrt{2\pi\sigma^2}} e^{-\frac{(x-\mu)^2}{2\sigma^2}}$  for which  $f'(x) = -\frac{1}{\sqrt{2\pi\sigma^6}} e^{-\frac{(x-\mu)^2}{2\sigma^2}} (x-\mu)$  and  $f''(x) = \frac{1}{\sqrt{2\pi\sigma^6}} e^{-\frac{(x-\mu)^2}{2\sigma^2}} \left( \frac{(x-\mu)^2}{\sigma^2} - 1 \right)$ .

(a) What are the domain and intercepts of  $f$ ? What are the asymptotics at  $\pm\infty$ ? Are there any vertical asymptotes? What are the asymptotics there?

**Solution:** The function is defined for all  $x$  and is always positive. We have  $f(0) = \frac{1}{\sqrt{2\pi\sigma^2}}$ .

For large  $x$  the function will decay rapidly (morally like  $e^{-x^2/2\sigma^2}$  even if that's not the correct asymptotics), so we have the horizontal asymptote  $y = 0$  on both sides.


(b) What are the intervals of increase/decrease? The local and global extrema?

**Solution:** Since  $\frac{1}{\sqrt{2\pi\sigma^6}} e^{-\frac{(x-\mu)^2}{2\sigma^2}}$  is always positive,  $f'(x) > 0$  when  $x < \mu$  and  $f'(x) < 0$  when  $x > \mu$ . Thus  $f$  is increasing on  $(-\infty, \mu)$ , decreasing on  $(\mu, \infty)$  and has a local (and global) maximum at  $x = \mu$ .

(c) What are the intervals of concavity? Any inflection points?

**Solution:** Since  $\frac{1}{\sqrt{2\pi\sigma^{10}}} e^{-\frac{(x-\mu)^2}{2\sigma^2}}$  is always positive, the sign of  $f''(x)$  is the same as that of  $((x-\mu)^2 - \sigma^2)$ . In particular  $f''(x) > 0$  when  $|x-\mu| > \sigma$ , that is on  $(-\infty, \mu - \sigma) \cup (\mu + \sigma, \infty)$ . Conversely  $f''(x) < 0$  when  $|x-\mu| < \sigma$ , that is on  $(\mu - \sigma, \mu + \sigma)$ . Finally we see there are inflection points at  $\mu \pm \sigma$ .

(d) Sketch a graph of  $f(x)$ .



(5) (Final, December 2007)  $\star\star$  Let  $f(x) = x\sqrt{3-x}$ .

(a) Find its domain, intercepts, and asymptotics at the endpoints.

**Solution:** The function is defined for if  $3-x \geq 0$  that is for  $x \leq 3$ . It is positive if  $x > 0$ , so if  $0 < x < 3$  and negative if  $x < 0$ , and thus crosses the axis at  $x = 0$ . As  $x \rightarrow -\infty$  we have  $x\sqrt{3-x} \sim x\sqrt{-x} \sim -|x|^{3/2}$ . As  $x \rightarrow 3$  we have  $x \sim 3(3-x)^{1/2}$ .

(b) What are the intervals of increase/decrease? The local and global extrema?

**Solution:** We have  $f'(x) = \sqrt{3-x} - \frac{x}{2\sqrt{3-x}} = \frac{2(3-x)-x}{2\sqrt{3-x}} = \frac{6-3x}{2\sqrt{3-x}} = \frac{3}{2} \cdot \frac{2-x}{\sqrt{3-x}}$ . Since  $\frac{3}{2\sqrt{3-x}}$  is always positive, the sign of  $f'(x)$  is determined by  $2-x$ . Thus  $f'$  is increasing on  $x < 2$ , decreasing for  $2 < x < 3$  and has its unique local maximum at  $x = 2$ .

(c) Given  $f''(x) = \frac{3x-12}{4}(3-x)^{-3/2}$ , what are the intervals of concavity? Any inflection points?

**Solution:** We have  $f''(x) = \frac{3(x-4)}{4(3-x)^{-3/2}}$  is always positive. Now the domain of the function is  $x < 3$  so  $x-4 < -1 < 0$  on the entire domain and  $f''(x) < 0$  for all  $x$  – so the function is concave down and has no inflection points.

(d) Sketch a graph of  $f(x)$ .

