

Math 100:V02 – SOLUTIONS TO WORKSHEET 7
THE CHAIN RULE

1. THE CHAIN RULE

(1) We know $\frac{d}{dy} \sin y = \cos y$.

(a) Expand $\sin(y + h)$ to linear order in h . Write down the linear approximation to $\sin y$ about $y = a$.

Solution: $\sin(y + h) \approx \sin y + h \cos y$ and $\sin y \approx \sin a + (y - a) \cos a$.

(b) Now let $F(x) = \sin(3x)$. Expand $F(x + h)$ to linear order in h . What is the derivative of $\sin 3x$?

Solution: $F(x + h) = \sin(3(x + h)) = \sin(3x + 3h)$ so we use $y = 3x$ in the previous example to get

$$\begin{aligned} F(x + h) &= \sin(3(x + h)) \\ &= \sin(3x + 3h) \\ &\approx \sin(3x) + (3h) \cos(3x) \\ &= \sin(3x) + (3 \cos(3x))h \end{aligned}$$

so the derivative is $\boxed{3 \cos(3x)}$.

(2) Write each function as a composition and differentiate

(a) e^{3x}

Solution: This is $f(g(x))$ where $g(x) = 3x$ and $f(y) = e^y$. The derivative is thus

$$e^{3x} \cdot \frac{d(3x)}{dx} = 3e^{3x}.$$

(b) $\sqrt{2x + 1}$

Solution: This is $f(g(x))$ where $g(x) = 2x + 1$ and $f(y) = \sqrt{y}$. Thus

$$\frac{df(g(x))}{dx} = f'(g(x))g'(x) = \frac{1}{2\sqrt{g}} \cdot 2 = \frac{1}{\sqrt{2x + 1}}.$$

(c) (Final, 2015) $\sin(x^2)$

Solution: This is $f(g(x))$ where $g(x) = x^2$ and $f(y) = \sin y$. The derivative is then

$$\cos(x^2) \cdot 2x = 2x \cos(x^2).$$

(d) $(7x + \cos x)^n$.

Solution: This is $f(g(x))$ where $g(x) = 7x + \cos x$ and $f(y) = y^n$. The derivative is thus

$$n(7x + \cos x)^{n-1} \cdot (7 - \sin x).$$

(3) (Final, 2012) Let $f(x) = g(2 \sin x)$ where $g'(\sqrt{2}) = \sqrt{2}$. Find $f'(\frac{\pi}{4})$.

Solution: By the chain rule, $f'(x) = g'(2 \sin x) \cdot \frac{d}{dx}(2 \sin x) = 2g'(2 \sin x) \cos x$. In particular,

$$\begin{aligned} f'(\frac{\pi}{4}) &= 2g'(2 \sin \frac{\pi}{4}) \cos \frac{\pi}{4} = 2g'(2 \frac{\sqrt{2}}{2}) \cdot \frac{\sqrt{2}}{2} \\ &= 2\sqrt{2} \cdot \frac{\sqrt{2}}{2} = 2. \end{aligned}$$

(4) Differentiate

(a) $7x + \cos(x^n)$

Solution: We apply linearity and then the chain rule:

$$\begin{aligned}\frac{d}{dx}(7x + \cos(x^n)) &= \frac{d(7x)}{dx} + \frac{d\cos(x^n)}{dx} \\ &= 7 + \frac{d\cos(x^n)}{d(x^n)} \cdot \frac{d(x^n)}{dx} \\ &= 7 - \sin(x^n) \cdot nx^{n-1}.\end{aligned}$$

(b) $e^{\sqrt{\cos x}}$

Solution: We repeatedly apply the chain rule:

$$\begin{aligned}\frac{d}{dx}e^{\sqrt{\cos x}} &= e^{\sqrt{\cos x}} \frac{d}{dx}\sqrt{\cos x} \\ &= e^{\sqrt{\cos x}} \frac{1}{2\sqrt{\cos x}} \frac{d}{dx}\cos x \\ &= -e^{\sqrt{\cos x}} \frac{\sin x}{2\sqrt{\cos x}}.\end{aligned}$$

(c) (Final 2012) $e^{(\sin x)^2}$

Solution: By the chain rule:

$$\begin{aligned}\frac{d}{dx}\left(e^{(\sin x)^2}\right) &= e^{(\sin x)^2} \frac{d}{dx}\left((\sin x)^2\right) \\ &= e^{(\sin x)^2} 2 \sin x \frac{d}{dx}\sin x \\ &= e^{(\sin x)^2} 2 \sin x \cos x \\ &= e^{(\sin x)^2} \sin(2x).\end{aligned}$$

(5) Suppose f, g are differentiable functions with $f(g(x)) = x^3$. Suppose that $f'(g(4)) = 5$. Find $g'(4)$.

Solution: Applying the chain rule we have $f'(g(x)) \cdot g'(x) = 3x^2$. Plugging in $x = 4$ we get $5g'(4) = 3 \cdot 4^2$ and hence $g'(4) = \frac{48}{5}$.

2. DIFFERENTIATING LOGARITHMS

(6) $\log(e^{10}) = \log(2^{100}) =$

Solution: $\log e^{10} = 10$ while $\log(2^{100}) = 100 \log 2$.

(7) Differentiate

(a) $\frac{d(\log(ax))}{dx} = \frac{d}{dt} \log(t^2 + 3t) =$

Solution: By the chain rule, the derivatives are: $\frac{1}{ax} \cdot a = \frac{1}{x}$ and $\frac{1}{t^2+3t} \cdot (2t+3) = \frac{2t+t}{t^2+3t}$. We can also use the logarithm laws first: $\log(ax) = \log a + \log x$ so $\frac{d}{dx}(\log ax) = \frac{d}{dx}(\log a) + \frac{d}{dx}(\log x) = \frac{1}{x}$ since $\log a$ is constant if a is. Similarly, $\log(t^2+3t) = \log t + \log(t+3)$ so its derivative is $\frac{1}{t} + \frac{1}{t+3}$.

(b) $\frac{d}{dx}x^2 \log(1+x^2) = \frac{d}{dr} \frac{1}{\log(2+\sin r)} =$

Solution: Applying the product rule and then the chain rule we get: $\frac{d}{dx}(x^2 \log(1+x^2)) = 2x \log(1+x^2) + x^2 \frac{1}{1+x^2} \cdot 2x = 2x \log(1+x^2) + \frac{2x^3}{1+x^2}$. Using the quotient rule and the chain rule we get

$$\frac{d}{dr} \frac{1}{\log(2+\sin r)} = -\frac{1}{\log^2(2+\sin r)} \cdot \frac{1}{2+\sin r} \cdot \cos r = -\frac{\cos r}{(2+\sin r)\log^2(2+\sin r)}.$$

(8) (Logarithmic differentiation) differentiate

$$y = (x^2 + 1) \cdot \sin x \cdot \frac{1}{\sqrt{x^3+3}} \cdot e^{\cos x}.$$

Solution: We have

$$\begin{aligned}\log y &= \log(x^2 + 1) + \log(\sin x) + \log\left(\frac{1}{\sqrt{x^3 + 3}}\right) + \log(e^{\cos x}) \\ &= \log(x^2 + 1) + \log(\sin x) - \frac{1}{2}\log(x^3 + 3) + \cos x.\end{aligned}$$

Differentiating with respect to x gives:

$$\frac{y'}{y} = \frac{2x}{x^2 + 1} + \frac{\cos x}{\sin x} - \frac{1}{2} \frac{3x^2}{x^3 + 3} - \sin x$$

and solving for y' finally gives

$$y' = \left(\frac{2x}{x^2 + 1} + \frac{\cos x}{\sin x} - \frac{3x}{2(x^3 + 3)} - \sin x \right) \cdot (x^2 + 1) \cdot \sin x \cdot \frac{1}{\sqrt{x^3 + 3}} \cdot e^{\cos x}.$$

(9) Differentiate using $f' = f \times (\log f)'$

(a) x^n

Solution: If $y = x^n$ then $\log y = n \log x$. Differentiating with respect to x gives $\frac{1}{y}y' = \frac{n}{x}$ so $y' = y \frac{n}{x} = nx^{n-1}$.

Solution: By the rule, $\frac{d}{dx}(x^n) = x^n \frac{d}{dx}(\log(x^n)) = x^n \left(\frac{n}{x}\right) = nx^{n-1}$.

(b) x^x

Solution: If $y = x^x$ then $\log y = x \log x$. Differentiating with respect to x gives $\frac{1}{y}y' = \log x + x \cdot \frac{1}{x} = \log x + 1$ so $y' = y(\log x + 1) = x^x(\log x + 1)$.

Solution: By the rule, $\frac{d}{dx}(x^x) = x^x \frac{d}{dx}(\log(x^x)) = x^x(\log x + 1)$.

Solution: We have $x^x = (e^{\log x})^x = e^{x \log x}$. Applying the chain rule we now get $(x^x)' = e^{x \log x}(\log x + 1) = x^x(\log x + 1)$.

(c) $(\log x)^{\cos x}$

Solution: By the logarithmic differentiation rule we have

$$\begin{aligned}\frac{d}{dx}(\log x)^{\cos x} &= (\log x)^{\cos x} \cdot \frac{d}{dx}(\cos x \log(\log x)) \\ &= -\sin x \log \log x (\log x)^{\cos x} + (\log x)^{\cos x} \cos x \frac{1}{\log x} \frac{1}{x} \\ &= -\sin x \log \log x (\log x)^{\cos x} + \cos x (\log x)^{\cos x - 1} \frac{1}{x}.\end{aligned}$$

(d) (Final, 2014) Let $y = x^{\log x}$. Find $\frac{dy}{dx}$ in terms of x only.

Solution: By the logarithmic differentiation rule we have

$$\begin{aligned}\frac{dy}{dx} &= y \frac{d \log y}{dx} = x^{\log x} \frac{d}{dx}(\log x \cdot \log x) \\ &= x^{\log x} \left(2 \log x \cdot \frac{1}{x} \right) = 2 \log x \cdot x^{\log x - 1}.\end{aligned}$$

3. MORE PROBLEMS

(10) Let $f(x) = g(x)^{h(x)}$. Find a formula for f' in terms of g' and h' .

Solution: By the logarithmic differentiation rule we have

$$\begin{aligned}f' &= f \cdot (h \log g)' \\ &= f \left(h' \log g + \frac{h}{g} g' \right) \\ &= h \cdot g^{h-1} \cdot g' + g^h \log g \cdot h' .\end{aligned}$$

Observe that this is the sum of what we'd get by applying the power law rule and the exponential rule.

- (11) Let $f(\theta) = \sin^2 \theta + \cos^2 \theta$. Find $\frac{df}{d\theta}$ without using trigonometric identities. Evaluate $f(0)$ and conclude that $\sin^2 \theta + \cos^2 \theta = 1$ for all θ .

Solution: By the chain rule $\frac{d}{d\theta} (\sin \theta)^2 = 2 \sin \theta \cos \theta$ and $\frac{d}{d\theta} (\cos \theta)^2 = 2 \cos \theta (-\sin \theta)$ so

$$\frac{df}{d\theta} = 2 \sin \theta \cos \theta - 2 \sin \theta \cos \theta = 0,$$

It follows that f is constant; since $f(0) = (\sin 0)^2 + (\cos 0)^2 = 1$ we have $f(\theta) = 1$ for all θ , which is the claim.

- (12) (“Inverse function rule”) suppose $f(g(x)) = x$ for all x .

(a) Show that $f'(g(x)) = \frac{1}{g'(x)}$.

Solution: Applying the chain rule we have $f'(g(x)) \cdot g'(x) = 1$.

(b) Suppose $g(x) = e^x$, $f(y) = \log y$. Show that $f(g(x)) = x$ and conclude that $(\log y)' = \frac{1}{y}$.

Solution: $f(g(x)) = \log(e^x) = x$. We then have $f'(e^x) = \frac{1}{g'(x)} = \frac{1}{e^x}$ so $f'(y) = \frac{1}{y}$ for all $y > 0$.

(c) Suppose $g(\theta) = \sin \theta$, $f(x) = \arcsin x$ so that $f(g(\theta)) = \theta$. Show that $f'(x) = \frac{1}{\sqrt{1-x^2}}$.

Solution: We have $f'(\sin \theta) = \frac{1}{\cos \theta} = \frac{1}{\sqrt{1-\sin^2 \theta}}$ so $f'(x) = \frac{1}{\sqrt{1-x^2}}$ for $-1 < x < 1$.

- (13) (Final, 2015) Let $xy^2 + x^2y = 2$. Find $\frac{dy}{dx}$ at the point $(1, 1)$.

Solution: Differentiating with respect to x we find $y^2 + 2xy \frac{dy}{dx} + 2xy + x^2 \frac{dy}{dx} = 0$ along the curve. Setting $x = y = 1$ we find that, at the indicated point,

$$3 + 3 \frac{dy}{dx} \Big|_{(1,1)} = 0$$

so

$$\frac{dy}{dx} \Big|_{(1,1)} = -1.$$